

# MUSCUREL®

thiocolchicoside

**Effective treatment**  
for **painful muscle contractures**  
in acute pathology  
(as from 16 y of age)

excellent efficacy  
low risk of side effects

## For more flexibility without pain

The patients is able to function normally again  
and to carry on the daily activities

### Medullary action without acting centrally

- no effect on respiration
- no risk of hypotension
- no harmful effects on kidney / liver functions
- no risk of addiction (as with benzos)
- no sedative effects
- no drug interactions



### POSOLOGY :

The recommended and maximum dose is 8 mg every 12 hours (or 16 mg a day), preferably after a meal.

The maximum duration of treatment is limited to 7 consecutive days.

Do not exceed the recommended dose or treatment duration.

**Not indicated below age 16.**

**Do not use in pregnant or breastfeeding women.**

The only thiocolchicoside available in 8 mg

Fewer doses

Very advantageous price



SCAN

For information about our other products,  
please visit [www.dafrapharma.com](http://www.dafrapharma.com)  
and create your personal account on the site.

# MUSCUREL®

thiocolchicoside



**Untangle the knots in the muscles  
when patients are shackled by pain**

For more flexibility without pain



Effective treatment for  
**PAINFUL MUSCLE CONTRACTURES**

(IN ACUTE SPINAL PATHOLOGY)

(from age 16)



**Setting the standard**

[www.dafrapharma.com](http://www.dafrapharma.com)

# MUSCUREL®

thiocolchicoside

## MUSCUREL®

- eliminates or considerably attenuates contracture of central origin
- reduces the passive resistance of the muscle to stretching
- reduces or eliminates residual contracture

*Thiocolchicoside is free of curariform effects and does not paralyze the muscle. The muscle relaxant effect is present thanks to the activation of GABA-A receptor antagonists and an agonist action on glycinergic receptors located in the brainstem and spinal cord.*

## MUSCUREL® :

- **no impairment of voluntary mobility**
- **eliminates any respiratory risk**
- **no influence on the cardiovascular system**

## Very good efficacy and utility in various cases

- no impact on the function of young people and workers
- easy to use - maximum 2 tablets per 24 h
- no respiratory side effects
- no risk of hypotension
- no effects/suppression in the brain
- no kidney/liver side effects
- no addiction

## DO NOT USE

- in pregnant or breastfeeding women
- in the event of an allergy to colchicine or one of the components of the medicinal product

## INDICATION

**MUSCUREL®** is used in adults and adolescents above age 16, as an adjuvant treatment for painful muscle contractures. It is used for acute painful conditions related to the spine.



### Painful muscle contractures with a spinal origin

after athletic overtraining - after excessive muscle use - overwork  
during a trauma - a contusion - a strain  
lumbago - stiff neck

## UNTANGLE THE KNOTS IN THE MUSCLES WHEN PATIENTS ARE SHACKLED BY PAIN

FOR MORE FLEXIBILITY WITHOUT PAIN

**Possible side effects:** digestive disorders - stomach pains. In case of diarrhea, dosage may be readjusted. In the event of a convulsive seizure, treatment should be stopped.

**Driving and use of machinery:** This drug may cause drowsiness. This risk increases if alcohol is consumed during treatment.

**Caution:** Do not exceed the maximum recommended dose (16 mg/day) or the maximum treatment duration of 7 consecutive days. Higher than recommended doses or long-term use should always be avoided.

**Muscurel®** is a medicinal product dispensed by prescription.



SCAN

For the full SMPC, please scan this QR code.