

Rhino PAROL[®]

paracetamol, chlorpheniramine maleate
& phenylephrine hydrochloride

**GET RID OF A STUFFY OR RUNNY NOSE
PUT AN END TO CONGESTION AND SNEEZING
PUT A STOP TO PAIN AND FEVER**



Usual dose: (as from 12 years of age)
1 tablet every 6 hours
max 4 tablets/day

CORYZA – FLU – COLD – RHINITIS – PHARYNGITIS –
SINUSITIS – BRONCHITIS – LARYNGITIS – ALLERGIES

For information about our other products,
please visit www.dafrapharma.com
and create your personal account.



Rhino PAROL[®]

paracetamol, chlorpheniramine maleate
& phenylephrine hydrochloride



MULTI-ACTION AGAINST COLD AND ALLERGY SYMPTOMS:

FROM 12 YEARS OF AGE

- Runny nose
- Stuffy nose
- Sinus congestion
- Sneezing
- Pain
- Fever



Setting the standard

www.dafrapharma.com

Rhino PAROL®

paracetamol, chlorpheniramine maleate
& phenylephrine hydrochloride



HIGHLY CONCENTRATED ACTIVE SUBSTANCES FOR A RAPID AND EFFECTIVE RELIEF

Paracetamol	650 mg	PAIN FEVER
Chlorpheniramine	4 mg	SNEEZING RUNNY NOSE ITCHY NOSE WATERY EYES
Phenylephrine	10 mg	SINUS CONGESTION STUFFY NOSE

Excellent efficacy

Start of the effect : after a few minutes

Duration of the effect : on average 4 to 6 hours

The use of RHINOPAROL® is not recommended during pregnancy or in nursing mothers.

In some patients who are receiving chlorpheniramine side effects such as drowsiness, dizziness, blurred vision, tremor and excitability may occur. Patients should be warned not to engage in hazardous activities while experiencing these side effects.

CORYZA

FLU

RHINITIS

COLD

PHARYNGITIS

SINUSITIS

BRONCHITIS

LARYNGITIS



ALLERGIES

RHINOSINUSITIS

GET RID OF A STUFFY OR RUNNY NOSE

PUT AN END TO CONGESTION AND SNEEZING

STOP SUFFERING FROM PAIN AND FEVER

CONTRA-INDICATIONS: RHINOPAROL® is contra-indicated in patients with a history of hypersensitivity to any of its active or inactive ingredients. It is also contra-indicated in benign prostatic hyperplasia, bladder neck obstruction, angle closure glaucoma, acute asthma, hyperthyroidism, severe hypertension, cardiac arrhythmias and cerebrovascular disorders.

UNDESIRABLE EFFECTS: Drowsiness, palpitation, blurred vision, dry mouth, anorexia, irritability, difficult urination, skin allergy, perspiration, tinnitus, tachycardia, photosensitivity, burning in the stomach.



For the full SMPC, please scan this QR code.

SCAN