

<b>AMIFER® SYRUP</b>	<b>AMIFER® JUNIOR SYRUP</b>	<b>AMIFER® FORTE TABLETS</b>																		
200 ml bottle with, per 5 ml: 30 mg elemental iron 2.50 mg folic acid 0.02 mg vitamin B12 Orange, mint and vanilla flavour	150 ml bottle with, per 5 ml: 50 mg elemental iron Cream flavour	Box of 30 tablets, per tablet: 100 mg elemental iron 0.350 mg folic acid																		
Prevention and treatment of iron deficiencies of various origins Iron deficiency, diet too low in iron Correction / treatment of iron deficiency anemia (haemoglobin level below normal)																				
For women who are preparing to conceive or pregnant women  For postpartum iron- deficiency anemia  To prevent deficiency of iron, folic acid and vitamin B12	Newborns and children  Can also be used in adults	Adolescents and adults who prefer tablets																		
Iron deficiencies																				
<table border="1"> <thead> <tr> <th>Age brackets</th> <th>Daily quantities</th> </tr> </thead> <tbody> <tr> <td>0 - 13 years old</td> <td>5 ml</td> </tr> <tr> <td>Adolescents &amp; adults</td> <td>10 ml</td> </tr> <tr> <td>Pregnancy &amp; allaitement</td> <td>10 ml</td> </tr> </tbody> </table>	Age brackets	Daily quantities	0 - 13 years old	5 ml	Adolescents & adults	10 ml	Pregnancy & allaitement	10 ml	<table border="1"> <thead> <tr> <th>Age brackets</th> <th>Daily quantities</th> </tr> </thead> <tbody> <tr> <td>Newborns - 2 years old</td> <td>1/4 teaspoon</td> </tr> <tr> <td>2 - 5 years old</td> <td>1/4 teaspoon, 1 to 2 times</td> </tr> <tr> <td>6 - 12 years old</td> <td>1 teaspoon, 1 to 2 times</td> </tr> <tr> <td>Adolescents &amp; adults</td> <td>2 teaspoons</td> </tr> </tbody> </table>	Age brackets	Daily quantities	Newborns - 2 years old	1/4 teaspoon	2 - 5 years old	1/4 teaspoon, 1 to 2 times	6 - 12 years old	1 teaspoon, 1 to 2 times	Adolescents & adults	2 teaspoons	1 tablet 1 to 2 times a day
Age brackets	Daily quantities																			
0 - 13 years old	5 ml																			
Adolescents & adults	10 ml																			
Pregnancy & allaitement	10 ml																			
Age brackets	Daily quantities																			
Newborns - 2 years old	1/4 teaspoon																			
2 - 5 years old	1/4 teaspoon, 1 to 2 times																			
6 - 12 years old	1 teaspoon, 1 to 2 times																			
Adolescents & adults	2 teaspoons																			
<b>Anemia:</b> Double the dose																				
<p style="text-align: center;"><b>AMIFER® also exists in IV form</b></p> <p style="text-align: center;">Difficulties taking iron orally, gastrointestinal absorption problems For chronic nephropathy Patients on dialysis Patients with heart failure Serious postoperative bleeding When losses are greater than the oral absorption capacity When the patient's clinical situation necessitates rapid iron administration</p>																				

**Dafra Pharma International**  
Head Office  
Slachthuisstraat 30/7, 2300 Turnhout, Belgium

# AMIFER®



AMIFER FORTE tablets  
AMIFER JUNIOR syrup  
AMIFER syrup

Oral iron, suitable for all patients

## IRON HEALTH FOR ALL AGES

Prevention and treatment of iron deficiencies  
of various origins

Iron deficiency sets in well before iron-deficiency anemia

General fatigue that lasts, lack of energy

Loss of efficacy / performance, concentration problems

Loss of muscle strength

Sleep problems, headaches and dizziness

Feeling chilled, increased vulnerability to infections, inflammation

Premature births or low birth weight in children

CONSIDER IRON DEFICIENCY

(even without anemia)



# AMIFER®

<b>AMIFER® SYRUP</b>	<b>AMIFER® JUNIOR SYRUP</b>	<b>AMIFER® FORTE TABLETS</b>
200 ml bottle with, per 5 ml: 30 mg elemental iron 2.50 mg folic acid  0.02 mg vitamin B12 Orange, mint and vanilla flavour Measuring cup	150 ml bottle with, per 5 ml: 50 mg elemental iron  Contains sucrose and sorbitol Cream flavour  Teaspoon with indications (1.25 ml - 2.5 ml - 5 ml)	Box of 30 tablets, per tablet: 100 mg elemental iron 0.350 mg folic acid
<b>Prevention</b> of iron and folic acid deficiencies. This syrup is indicated for <b>people with iron deficiency or whose diet is low in iron</b> , during pregnancy, after childbirth and <b>prevention</b> of anemia and iron and folic acid deficiency.	In the <b>treatment and prevention</b> of iron <b>deficiencies of various origins</b> and for people with iron-deficiency <b>anemia</b> , or to treat iron deficiency during <b>childhood</b> or during <b>pregnancy and breastfeeding</b> .	Use in the treatment of all iron <b>deficiencies</b> of various causes, in the <b>prevention</b> and treatment of iron and <b>folic acid</b> deficiencies during <b>pregnancy</b> and breastfeeding and in the treatment of <b>iron-deficiency anemia</b> .

## Do not take oral forms of AMIFER® (consult the SMPCs for complete information)

- If you are hypersensitive (allergic) to iron or one of its components.
- For iron accumulation (iron overload) (hemochromatosis, hemosiderosis, chronic hemolysis) and problems inherent in iron consumption (sideroblastic anemias, lead poisoning anemias, thalassemias).
- For non-iron-deficiency anemia
- For conditions that need to receive frequent and continuous blood transfusions,
- For megaloblastic anemia resulting from an isolated vitamin B12 deficiency or isolated folic acid deficiency.

## PREGNANCY AND BREASTFEEDING

Oral forms of **AMIFER®** can be administered safely during pregnancy and breastfeeding if the physician or pharmacist recommends its use.

## DRIVING AND USING MACHINES

No adverse effects related to driving or using machines have been observed.

## OVERDOSE

In the event of overdose, immediately contact your doctor or pharmacist.

When taking oral iron formulations, the colour of stool may darken; this is a normal phenomenon that does not require any special measures. It will not induce false positive tests when checking the presence of blood in the stool, so it is not necessary to interrupt treatment during this test.

## POSSIBLE SIDE EFFECTS

Allergic reactions, indigestion, abdominal pain, nausea or vomiting, burning sensation in the stomach, constipation, diarrhea, bitter taste in the mouth, mild abdominal pain, itchy blisters on the skin, rash, redness, headache, change in the color of urine and stools.

## STORAGE

Store at a temperature below 30°C. Do not use this medicinal product if the expiry date (Exp.) on the packaging or bottle is exceeded.

# THE BENEFITS OF ORAL FORMS OF AMIFER®

BEFORE (possible symptoms of iron deficiencies)	AFTER
Mental fatigue Concentration difficulties	Calm and serenity regained Able to concentrate again Regain drive ...
Physical fatigue Lack of strength and performance	Be able to do physical activities again Regain drive
Concentration problems Reduction of cognitive functions	Do mental work again, study, school performance, etc
Sleep disorders	Sleep through the night without awakening Awake refreshed
Frequent headaches	Fewer headaches
Feeling chilled	Better withstand cold
Increased vulnerability to infections	Fewer illnesses: good resistance
Hair loss Weak and ridged nails Chapped lips	Better physical condition
Premature birth Weight deficit	Better chance to have a pregnancy with no problems...



## IRON HEALTH FOR ALL AGES