

Amlovie®

amlodipine

POSODOLOGY

The initial recommended dosage is 5 mg once daily.

This dose may be increased up to 10 mg once daily.

Adjust the dosage according to the blood pressure goals. In general, wait 7 to 14 days between the titration steps. However, if the clinical situation justifies it, it is possible to titrate more quickly, as long as the patient is frequently evaluated.

It is preferable to take **AMLOVIE®** every day at the same time with a glass of water.

Reducing the blood pressure reduces the risk of fatal and non-fatal cardiovascular events, mainly cerebral vascular accidents and myocardial infarctions.

Take your patients to heart

Relax the vessels...

... to reduce blood pressure

... to improve blood supply to the heart muscle

DAFRA PHARMA INTERNATIONAL
Head Office – Slachthuisstraat 30/7 – 2300 Turnhout – Belgium

3FIAMLOE-2021



Setting the standard
www.dafrapharma.com

Amlovie®


amlodipine



**Blood pressure
Angina**



The gold standard calcium antagonist



Relax the vessels

... to reduce blood pressure
... to improve blood supply to the heart muscle

High-quality European calcium antagonist
AT A VERY ACCESSIBLE PRICE



Setting the standard

www.dafrapharma.com

Amlovie®

amlodipine

AMLOVIE® is used to treat **high pressure** or **angina**.

- In patients with high blood pressure values, **AMLOVIE®** acts by relaxing the blood vessels, so that the blood passes through them more easily.
- In patients with angina, **AMLOVIE®** acts by improving blood supply to the heart muscle, which therefore receives more oxygen, preventing the onset of chest pain.

How to take Amlovie® ?

- The initial recommended dosage is 5 mg once daily.
- This dose may be increased up to 10 mg once daily.
- It is preferable to take **AMLOVIE®** at the same time every day with a glass of water.

Special populations:

- Elderly : **AMLOVIE®** shows equivalent good tolerability in elderly patients and younger patients. Normal dosage regimens are recommended in the elderly, but increasing the dosage should be done with caution.
- Hepatic impairment The dose should be determined carefully and treatment must start with a dose located at the lower limit of the therapeutic range. Initiate treatment with **AMLOVIE®** at the lowest possible dose and perform a slow titration in patients with severe hepatic impairment
- Renal impairment : The usual dosage is recommended.
- Paediatric population : Children and adolescents aged 6 to 17 years with high blood pressure: the recommended oral high blood pressure treatment dosage in children aged 6 to 17 years is 2.5 mg, once daily as the initial dose. This dose may be increased up to 5 mg once daily if the desired blood pressure is not reached after 4 weeks. Dosages above 5 mg once daily have not been studied in paediatric patients. An amlodipine dosage of 2,5 mg is not possible with **AMLOVIE®** 10 mg. Children below age 6 years: There are no available data.

In patients with high blood pressure, **AMLOVIE®** can be used in combination with a thiazide diuretic, an alpha blocker, a beta blocker or an angiotensin conversion enzyme inhibitor.

In angina, **AMLOVIE®** can be used as a monotherapy or in combination with other angina treatments in patients with angina resistant to nitrate derivatives and/or appropriate doses of beta blockers.

Side effects: The most commonly reported side effects in treatment are: drowsiness, dizziness, headache, palpitations, flushing, abdominal pain, nausea, ankle swelling, oedema and fatigue.

Do not take **AMLOVIE®** in cases of allergy to amlodipine or other calcium channel blockers or any of the ingredients of this medication, or low blood pressure, aortic valve stenosis or cardiogenic shock and heart failure after a heart attack.

VASOCONSTRICTION

increase in blood pressure
HIGH BLOOD PRESSURE :
a silent killer

reduction of distal blood flow
and oxygenation
ANGINA:
a harbinger of a heart attack

Amlovie®



**AMLOVIE® fights
high blood pressure**

**AMLOVIE® treats and/or
prevents angina**

Take your patients to heart

Relax the vessels...

... to reduce blood pressure

... to improve blood supply to the heart muscle

AMLOVIE® is a medicinal product dispensed by prescription.

For the complete SMPC, consult www.dafrapharma.com