



Don't waste time

Rely on Daфра Pharma

We deliver a wide range of high quality
pharmaceutical products and services which are at
the reach of every patient

www.dafrapharma.com

Dafra Pharma's

Third-generation cephalosporine

Cetafor®
500 mg IV

Powder and solvent
for 5 ml IV injection



Cetafor®
1000 mg IV

Powder and solvent
for 10 ml IV injection



Your reliable partner in the treatment of infectious diseases

Cetafor® IV



DOSAGE

Adults: 1 - 2 g per day*
Children: 50 - 75 mg/kg/day*
100 mg/kg/day*
in case of meningitis

* Do not exceed 4 grams daily

PREPARATION FOR INJECTIONS:

- dissolve 500 mg **Cetafor® IV** in 5 ml water for injection
- dissolve 1000 mg **Cetafor® IV** in 10 ml water for injection
- administer the IV injection over 2 - 4 minutes

Setting the Standard

www.dafrapharma.com

For an extensive scientific product information,
we refer to the leaflet in the packaging

Cetafor® IV

LOWER RESPIRATORY TRACT INFECTIONS caused by *Streptococcus pneumoniae*, *Staphylococcus aureus*, *Haemophilus influenzae*, *Haemophilus parainfluenzae*, *Klebsiella pneumoniae*, *Escherichia coli*, *Enterobacter aerogenes*, *Proteus mirabilis* or *Serratia marcescens*.

ACUTE BACTERIAL OTITIS MEDIA caused by *Streptococcus pneumoniae*, *Haemophilus influenzae* (including beta-lactamase producing strains) or *Moraxella catarrhalis* (including beta-lactamase producing strains).

SKIN AND SKIN STRUCTURE INFECTIONS caused by *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Streptococcus pyogenes*, Viridans group streptococci, *Escherichia coli*, *Enterobacter cloacae*, *Klebsiella oxytoca*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Morganella morganii*, *Pseudomonas aeruginosa*, *Serratia marcescens*, *Acinetobacter calcoaceticus*, *Bacteroides fragilis* or Peptostreptococcus species.

URINARY TRACT INFECTIONS (complicated and uncomplicated) caused by *Escherichia coli*, *Proteus mirabilis*, *Proteus vulgaris*, *Morganella morganii* or *Klebsiella pneumoniae*.

UNCOMPLICATED GONORRHEA (cervical/urethral and rectal) caused by *Neisseria gonorrhoeae*, including both penicillinase- and nonpenicillinase-producing strains, and pharyngeal gonorrhoea caused by nonpenicillinase-producing strains of *Neisseria gonorrhoeae*.

PELVIC INFLAMMATORY DISEASE caused by *Neisseria gonorrhoeae*.

BACTERIAL SEPTICEMIA caused by *Staphylococcus aureus*, *Streptococcus pneumoniae*, *Escherichia coli*, *Haemophilus influenzae* or *Klebsiella pneumoniae*.

BONE AND JOINT INFECTIONS caused by *Staphylococcus aureus*, *Streptococcus pneumoniae*, *Escherichia coli*, *Proteus mirabilis*, *Klebsiella pneumoniae* or *Enterobacter* species.

INTRA-ABDOMINAL INFECTIONS caused by *Escherichia coli*, *Klebsiella pneumoniae*, *Bacteroides fragilis*, *Clostridium* species or Peptostreptococcus species.

MENINGITIS caused by *Haemophilus influenzae*, *Neisseria meningitidis* or *Streptococcus pneumoniae*. Ceftriaxone has also been used successfully in a limited number of cases of meningitis and shunt infection caused by *Staphylococcus epidermidis* and *Escherichia coli*.

SURGICAL PROPHYLAXIS: preoperative administration of a single 1 g dose

Setting the Standard

www.dafrapharma.com

For an extensive scientific product information,
we refer to the leaflet in the packaging