

# MUSCUREL®

thiocolchicoside

**Effective adjuvant treatment (from age 16) for**

## **PAINFUL MUSCLE CONTRACTURE** (IN ACUTE SPINAL PATHOLOGY)

excellent effectiveness with low risk of side effects

**The patient can continue to freely go about  
their daily activities.**

### **Medullary action without acting centrally**

- no effect on respiration
- no risk of hypotension
- no harmful effects on kidney / liver functions
- no risk of addiction (as with benzos)
- no sedative effects
- no drug interactions



### **POSODOLOGY :**

The recommended and maximum dose is 8 mg every 12 hours (or 16 mg a day), preferably after a meal.

The maximum duration of treatment is limited to 7 consecutive days.

Do not exceed the recommended dose or treatment duration.

**Not indicated below age 16.**

**Do not use in pregnant or breastfeeding women.**

The only thiocolchicoside available in 8 mg  
Fewer doses  
Very advantageous price

**DAFRA PHARMA INTERNATIONAL**  
Head Office – Slachthuisstraat 30/7 – 2300 Turnhout – Belgium

3FIMPRELE 2021

# MUSCUREL®

thiocolchicoside



## Loosen knots in muscles when pain binds you

For more flexibility without pain



Effective treatment for  
**PAINFUL MUSCLE CONTRACTURE**  
(IN ACUTE SPINAL PATHOLOGY)  
(from age 16)



**Setting the standard**

[www.dafrapharma.com](http://www.dafrapharma.com)

# MUSCUREL®

thiocolchicoside

## **MUSCUREL®**

- eliminates or considerably attenuates contracture of central origin
- reduces the passive resistance of the muscle to stretching
- reduces or eliminates residual contracture

*Thiocolchicoside is free of curariform effects and does not paralyze the muscle. The muscle relaxant effect is present thanks to the activation of GABA-A receptor antagonists and an agonist action on glycinergic receptors located in the brainstem and spinal cord.*

## **MUSCUREL® :**

- **no impairment of voluntary mobility**
- **eliminates any respiratory risk**
- **no influence on the cardiovascular system**

## **Very good efficacy and utility in various cases**

- no impact on the function of young people and workers
- easy to use - maximum 2 tablets per 24 h
- no respiratory side effects
- no risk of hypotension
- no effects/suppression in the brain
- no kidney/liver side effects
- no addiction

## **DO NOT USE**

- in pregnant or breastfeeding women
- in the event of an allergy to colchicine or one of the components of the medicinal product

## INDICATION

**MUSCUREL®** is used in adults and adolescents above age 16, as an adjuvant treatment for painful muscle contractures. It is used for acute painful conditions related to the spine.



### Painful muscle contractures with a spinal origin

after athletic overtraining - after excessive muscle use - overwork  
during a trauma - a contusion - a strain  
lumbago - stiff neck

## LOOSEN KNOTS IN MUSCLES WHEN PAIN BINDS YOU

**Possible side effects :** digestive problems - stomach problems. In the event of diarrhea, the dosage may be readjusted. In the event of a seizure, the treatment must be discontinued. In the event of liver damage during treatment with this medicinal product (for example: loss of appetite, nausea, vomiting, abdominal discomfort, fatigue, dark urine, jaundice, pruritus)

**Fertility :** This medicinal product may lead to male fertility problems by potential alteration of the sperm cells (abnormal number of chromosomes); this has been demonstrated in laboratory studies.

**Driving and using machines :** This medicinal product can commonly cause drowsiness. You should take this into account before driving or using machines. Drowsiness increases if you consume alcohol or medicines containing alcohol during treatment.

Muscurel® is a medicinal product dispensed by prescription.  
For the complete SMPC, consult [www.dafrapharma.com](http://www.dafrapharma.com)